

EXact-Cut™ NotI Restriction Endonuclease

Catalog Number: EXNA027

Size: 500 Units

For Research Use Only. Not Intended for Diagnostic or Therapeutic Use.

EXREprotein™

More information: info@exreprotein.com

Product Details

| | | |
|--|--|---------------------------|
| Description | EXact-Cut™ NotI Restriction Endonuclease is engineered for high specificity, reduced star activity, and time saving DNA digestion in 5-15 minutes. To simplify experimental design using multiple restriction enzymes, our entire range of EXact-Cut™ restriction endonucleases are 100% active in our EXact-Cut™ buffer (included) and are optimized for single-tube reactions along with digestion and ligation protocols. | |
| Restriction Enzyme Site | 5'...GC↓GGCC GC...3' 3'...CG CCGG↑CG...5' Isoschizomers: CciNI (Isoschizomers may have different methylation sensitivities) | |
| Unit Definition | One unit is defined as the amount of enzyme required to digest 1 µg of λ DNA in 1 hour at 37°C in a total reaction volume of 50 µL. | |
| Recommended Reaction Conditions | 1 x EXact-Cut™ Buffer Incubate at 37°C Refer to Protocol for reaction setup | |
| Heat Inactivation | 1. Incubate at 80°C for 20 minutes 2. Add appropriate volume of 6X Gel Loading Dye, according to the reaction system | |
| Components | EXact-Cut™ NotI (10 Units/µL) EXact-Cut™ 10X Buffer 6X Gel Loading Dye, Purple | 500 Units 1 mL 1 mL |

Storage and Preparation

| | |
|------------------------------|-------------------------------------|
| Shipping | Shipped on blue ice. |
| Stability and Storage | Store at -20°C for up to 24 months. |

Protocol

Protocol for Rapid DNA Digestion

1. Add the following components on ice in the indicated order:

| | Plasmid DNA | PCR Product | Genomic DNA |
|--|-------------|-------------|-------------|
| DNA | ≤ 1 µg | ≤ 0.2 µg | ≤ 5 µg |
| EXact-Cut™ 10X Buffer | 2 µL | 3 µL | 5 µL |
| ddH ₂ O, make up to final volume indicated: | 20 µL | 30 µL | 50 µL |
| EXact-Cut™ NotI | 10 Units | 10 Units | 30-50 Units |

Note: DNA should be free of phenol, chloroform, ethanol, EDTA, detergents or high concentrations of salts. For compatibility with other common buffers, see the chart on page 2.

- Gently mix or flick the tube to mix (do not vortex), then immediately follow with a quick spin-down in a microcentrifuge.
- Incubate at 37°C for the indicated sample type: plasmid DNA (15 minutes), PCR product (15-30 minutes), or genomic DNA (30-60 minutes)
- Optional: inactivate the enzyme at 80°C for 20 minutes and add an appropriate amount of 6X Gel Loading Dye, according to the reaction system.

Protocol for Multiple Digestion of DNA

- Use 10 Units of each enzyme and scale up to the reaction conditions accordingly.
- The combined volume of the enzymes in the reaction mixture **should not** exceed **1/10** of the total reaction volume.
- If the enzymes require different reaction temperatures, start with the enzyme requiring the lowest temperature, followed by the next enzyme(s) and incubate at the higher temperature.

Note: For total reaction volumes > 20 µL, the incubation time should be increased accordingly in a water bath.

Number of Recognition Sites in DNA

| λ DNA | Φ X174 | pBR322 | pUC57 | pUC18/19 | SV40 | M13mp18/19 | Adeno2 |
|---------------|-------------|--------|-------|----------|------|------------|--------|
| 0 | 0 | 0 | 0 | 0 | 0 | 0 | 7 |

Methylation Effects on Digestion

| Dam | Dcm | CpG | EcoKI | EcoBI |
|-----------|-----------|----------|-----------|-----------|
| No effect | No effect | Impaired | No effect | No effect |

Activity in Common Buffers

| | EXact-Cut™ Buffer | Takara QuickCut™ Buffer | Thermo Scientific FastDigest Buffer | NEB CutSmart® Buffer |
|----------|----------------------|----------------------------|--|-------------------------|
| Activity | 100% | 100% | 100% | 100% |

Application Notes

Functional Test

A 20 μ L reaction in EXact-Cut Buffer containing 1 μ g of p615 DNA and 10 Units of EXact-Cut NotI incubated for 15 minutes at 37°C results in complete digestion as determined by agarose gel electrophoresis.

Digestion-Ligation Test

At the optimal reaction temperature, the DNA was digested using 10 Units of EXact-Cut NotI and the digestion product was recovered. The DNA fragments were ligated using an appropriate amount of T4 DNA Ligase at 22°C. After the ligation product was recovered, it was able to be recut with EXact-Cut NotI.

Non-Specific Endonuclease Activity Test

At the optimal reaction temperature, 10 Units of EXact-Cut NotI was incubated in 20 μ L reaction volume in EXact-Cut Buffer with 1 μ g of supercoiled plasmid DNA for 4 hours. Undigested, supercoiled plasmid DNA was detected using agarose gel electrophoresis.

Blue/White Screening Assay

An appropriate vector containing the *lacZ* gene was digested using 10 Units of EXact-Cut NotI. The digested product was ligated and transformed into *E. coli* cells plated on plates with X-Gal, IPTG and appropriate antibiotic. The successfully ligated *lacZ* gene expresses beta-galactosidase and gives rise to a blue colony, while an interrupted gene (due to degraded DNA end) gives rise to a white colony. Less than 1% white colonies were observed.